

Heavy Ion Initial Conditions and Correlations Between Higher Moments in the Spatial Anisotropy

J. L. Nagle¹ and M. P. McCumber¹

¹*University of Colorado at Boulder**

(Dated: May 3, 2011)

Fluctuations in the initial conditions for relativistic heavy ion collisions are proving to be crucial to understanding final state flow and jet quenching observables. The initial geometry has been parametrized in terms of moments in the spatial anisotropy (i.e. $\epsilon_2, \epsilon_3, \epsilon_4, \epsilon_5 \dots$), and it has been stated in multiple published articles that the vector directions of odd moments are uncorrelated with the even moments and the reaction plane angle. In this article, we demonstrate that this is incorrect and that a substantial non-zero correlation exists between the even and odd moments in peripheral Au+Au collisions. These correlations persist for all centralities, though at a very small level for the 0-55% most central collisions.

One proposal for modeling the initial geometry fluctuations in relativistic heavy ion collisions is utilizing a Monte Carlo Glauber calculation [1]. Using this model and the initial transverse positions of the struck nucleons, referred to as participants, one can calculate the participant eccentricity ϵ_2 and higher moments [2]. In [2], these moments are defined by:

$$\epsilon_n = \frac{\sqrt{\langle r^2 \cos(n\phi_{part}) \rangle^2 + \langle r^2 \sin(n\phi_{part}) \rangle^2}}{\langle r^2 \rangle} \quad (1)$$

where n is the n th moment of the spatial anisotropy calculated relative to the mean position. The major axis associated with the n th moment is defined by:

$$\psi_n = \frac{\text{atan2}(\langle r^2 \sin(n\phi_{part}) \rangle, \langle r^2 \cos(n\phi_{part}) \rangle)}{n} \quad (2)$$

We show an example event display in Figure 1 that includes the positions of the participant nucleons, and a visualization of the $\epsilon_2, \epsilon_3, \epsilon_4$, and ϵ_5 moments. We have drawn the vector direction of each along the long-axis of the associated moment. The n th moment has an n -multiplet of directions that are equally valid, separated by $2\pi/n$.

I. PREVIOUS RESULTS

In [2], the authors state that the minor axis of triangularity (i.e. ϵ_3) is found to be uncorrelated with the minor axis of eccentricity (i.e. ϵ_2) in Monte Carlo Glauber calculations. This conclusion is repeated in [3] where it is stated that the orientation of triangular overlap shape due to fluctuations is random relative to the event-plane direction as determined by the elliptic anisotropy. This conclusion is then generalized in [4] where these authors state that the angles for the odd and even harmonics are uncorrelated.

There have been many initial studies for how this spatial anisotropy translates into the final momentum space distribution of particles (e.g. [2, 5–9]). The fact that the angular orientations are uncorrelated between odd and even moments has an important impact on the methodology for experiments to determine these momentum anisotropy moments v_n and for two-particle correlation measurements, with relevance for jet quenching observables.




FIG. 1. (Color online) Monte Carlo Glauber event display for a sample Au+Au collision. The grey circles represent the positions of all nucleons. The green (red) circles are participant nucleons from the left (right) nucleus. The vector directions for the $n = 2 - 5$ and the spatial anisotropy pattern they represent are overlaid. These are centered on the mean position as indicated by a black square.

* jamie.nagle@colorado.edu




FIG. 2. (Color online) Monte Carlo Glauber Au+Au distribution for relative angular difference ($\epsilon_2 - \epsilon_3$) [radians] for different selections in N_{part} . A flat distribution at one from angular separations $0.0-\pi/6$ indicates totally uncorrelated quantities.

II. METHODOLOGY

We set out to confirm the findings of the above papers using the Monte Carlo Glauber framework. We have utilized the standard PHOBOS Monte Carlo Glauber code [10] with Woods-Saxon parameters and settings ($R_0 = 6.38$ fm, $a = 0.535$ fm, $d_{min} = 0$ fm). We show the angular distribution between ϵ_2 and ϵ_3 for a set of Au+Au number of participant (N_{part}) selections in Figure 2. Because of the two (three) fold symmetry for the ϵ_2 (ϵ_3) moments, if the two angles are uncorrelated, the distribution should be flat at one from $0.0 - \pi/6$ radians. A clear correlation is found between the two moments in peripheral Au+Au events. The correlation strength then decreases for more central events [11].

Previous studies have noted that correlations exist between the positions of participating nucleons and are intrinsic to all Monte Carlo Glauber calculations [12]. And at the same time, previous publications assumed that this would not lead to correlations in the vector direction of odd and event moments since the odd moments are purely the result of fluctuations.

We quantify the degree of correlation by taking the root-mean-square (RMS) of the distribution. If there were no correlation between these moments, the RMS of the angular separation would be 0.302 radians. The results as a function of N_{part} are shown in Figure 3. For our initial view, we have zoomed in for $N_{part} < 100$. The results confirm the strong correlation between the two angles (i.e. the large downward deviation in the RMS from the flat case) for $N_{part} \approx 10$, the deviation then weakens for larger N_{part} values. In this figure, the cor-




FIG. 3. (Color online) The root-mean-square (RMS) of the angular difference between ϵ_2 and ϵ_3 as a function of number of participant nucleons, N_{part} , for Au+Au, Cu+Cu, and Si+Si collisions, as well as the expectation of no correlation. We plot $N_{part} < 100$ to highlight the significant correlations that are present at small N_{part} .




FIG. 4. (Color online) The root-mean-square (RMS) of the angular difference between ϵ_2 and ϵ_3 as a function of number of participant nucleons, N_{part} , for Au+Au, Cu+Cu, and Si+Si collisions, as well as the expectation of no correlation. We zoom in on the vertical axis to highlight the small deviations from no correlation that remain at large N_{part} .

relation appears to disappear for $N_{part} > 50$. However, in Figure 4, we show the full N_{part} range zooming the vertical axis around the default value. We observe a very small remaining anti-correlation for $N_{part} \approx 80 - 150$, which translates into an approximate 1% lower probability of having the two moments within $0.0-\pi/12$ and a correspondingly higher probability of having the two moments separated by $\pi/12-\pi/6$. There is a similar magnitude positive correlation for $N_{part} > 200$.

For the lower N_{part} region where the alignment is strongest, the correlation may be due to small number fluctuations of the particular geometry in that impact pa-




FIG. 5. (Color online) The RMS angular difference between the reaction plane, Ψ_{RP} , and ϵ_3 as a function of N_{part} for Au+Au collisions. A weak correlation is found between the 3rd-order moment and the reaction plane direction for small N_{part} . Note the zoomed vertical scale.




FIG. 6. (Color online) The root-mean-square (RMS) of the angular difference between ϵ_2 and ϵ_5 as a function of number of participant nucleons, N_{part} , for Au+Au, Cu+Cu, and Si+Si collisions, as well as the expectation of no correlation. The result demonstrates that a similar behavior holds for correlation between ϵ_2 and higher order odd moments.

parameter range for Au+Au. Thus, also shown in Figure 3 are calculations from Si+Si and Cu+Cu collisions. One can see that the angular correlation largely tracks with N_{part} and thus it is the number fluctuations and not the average geometry that dominate the correlation. However, the *Si*, *Cu*, *Au* results do not scale perfectly versus N_{part} , and thus the particular geometric configuration plays some role, as one might expect since the geometry correlates with the magnitude of these moments.

Furthermore, the authors of [13] state that the fluctuations are random with respect to the reaction plane (the plane defined the line between the centers of the nuclei and the longitudinal axis). We have tested this as well




FIG. 7. (Color online) The root-mean-square (RMS) of the angular difference between ϵ_2 and ϵ_4 as a function of number of participant nucleons, N_{part} , for Au+Au, Cu+Cu, and Si+Si collisions, as well as the expectation of no correlation. The result shows the same procedure applied to the difference between the lowest even-order moments where a more significant correlation results from the average geometry.

and show the resulting RMS angular separation between the ϵ_3 and the reaction plane in Figure 5. We find a similar trend as shown above. The correlation between ϵ_3 and the reaction plane is also restricted to small numbers of participating nucleons. The correlation here is much weaker than what was found between ϵ_2 and ϵ_3 .

We also show the angular correlation between ϵ_2 and ϵ_5 in Figure 6. One again sees a strong correlation in peripheral collisions and then smaller correlations for larger numbers of participants. Finally, in Figure 7, we show the angular correlation between ϵ_2 and ϵ_4 . These two even moments are expected to be highly correlated because the initial overlap in mid-central collisions has an approximately elliptical shape, which can be well described by a combination of aligned ϵ_2 and ϵ_4 moments if it has a large enough eccentricity. This correlation tracks with the geometry (i.e. impact parameter), and so the resulting RMS values for Au+Au, Cu+Cu, and Si+Si do not track each other when plotted as a function of N_{part} .

In the case of nearly ideal hydrodynamics, the spatial eccentricities may be translated into directly measurable momentum anisotropies (v_n). The relative degree of correlation between v_2 and v_4 is of great interest because it directly impacts a possible difference in the observed v_4 determined via the event plane method (using the second moment to determine the plane) and a two-particle Fourier decomposition method [14]. The v_3 moment is identically zero by symmetry when measured with respect to the second moment event plane. However, the vector direction correlation can be determined via event-by-event separate second and third-moment event plane measurements. Such measurements would provide potentially sensitive tests of the fluctuations in the Monte Carlo Glauber geometry. Despite the effect being largest

in peripheral events where nearly inviscid hydrodynamics may not apply, it should prove fruitful to see if such angle correlations persist (which do not require a linear translation of ϵ_n moments into v_n momentum anisotropies).

III. SUMMARY

In summary, we find that contrary to previous publications, there is a definite non-zero correlation within the Monte Carlo Glauber calculation between the angular directions of ϵ_2 and ϵ_3 (and more generally between even and odd moments). A much weaker correlation between the reaction plane and ϵ_3 is also found. The effects are strongest for peripheral Au+Au events, and the measurement of such correlations may provide an interesting test of geometric fluctuations within the Monte Carlo Glauber

calculation. It will be instructive to measure these possible correlations and care must be taken to quote the exact sensitivity of such measures and extend them to the very most peripheral events. The multi-dimensional correlation between the magnitude of the eccentricity orders and their angular orientations may also prove important and the direct integration of Monte Carlo calculations of initial state geometries may be warranted in many studies to account for the full set of correlations.

ACKNOWLEDGMENTS

We thank B. Alver, P. Steinberg, and W. Zajc for valuable discussions. We acknowledge funding from the Division of Nuclear Physics of the U.S. Department of Energy under Grant No. DE-FG02-00ER41152.

-
- [1] M. L. Miller, K. Reygers, S. J. Sanders, and P. Steinberg, *Ann. Rev. Nucl. Part. Sci.* **57**, 205 (2007).
 - [2] B. Alver and G. Roland, *Phys.Rev.* **C81**, 054905 (2010).
 - [3] H. Agakishiev, M. Aggarwal, Z. Ahammed, A. Alakhverdyants, I. Alekseev, et al. (2010), arXiv:1010.0690.
 - [4] R. A. Lacey, R. Wei, N. Ajitanand, and A. Taranenko (2010), arXiv:1009.5230.
 - [5] G.-Y. Qin, H. Petersen, S. A. Bass, and B. Muller (2010), arXiv:1009.1847.
 - [6] B. H. Alver, C. Gombeaud, M. Luzum, and J.-Y. Ollitrault, *Phys.Rev.* **C82**, 034913 (2010).
 - [7] H. Holopainen, H. Niemi, and K. J. Eskola (2010), arXiv:1007.0368.
 - [8] D. Teaney and L. Yan (2010), arXiv:1010.1876.
 - [9] B. Schenke, S. Jeon, and C. Gale (2010), arXiv:1009.3244.
 - [10] B. Alver, M. Baker, C. Loizides, and P. Steinberg (2008), arXiv:0805.4411.
 - [11] B. Alver, private communication, further calculation confirms the ϵ_2 - ϵ_3 correlation in peripheral events. The original publication [2] only checked for $N_{part} > 60$.
 - [12] W. Broniowski, P. Bożek, and M. Rybczyński, *Phys. Rev. C* **76**, 054905 (2007).
 - [13] H. Petersen, G.-Y. Qin, S. A. Bass, and B. Muller (2010), arXiv:1008.0625.
 - [14] M. Luzum (2010), arXiv:1011.5773.